

JRC MARS Bulletin

Crop monitoring in Europe

May 2020

Reduced yield expectations at EU level

Positive outlook in Mediterranean region and northern Europe

At EU level, the yield forecasts for almost all crops were revised downwards. Most markedly for rapeseed, winter barley and winter wheat. In the case of spring barley, downward revisions of the yield forecasts in most countries were offset by an upward revision in Spain. For summer crops, the season just started and most of the yield forecasts reported at country level, are still based on historical trends or average values.


In large parts of western and central Europe, weather conditions were similar to those reported in the April Bulletin: predominantly above-average temperatures, high radiation levels, and insufficient precipitation to replenish soil moisture levels for the growth of winter and early-planted spring crops. In eastern Romania and western Ukraine, crops are stunted and present signs of wilting and early leaf senescence.

For late planted spring and summer crops, the concerns raised in the April Bulletin for these regions were mitigated. However, emerging crops often present poor stands due to the preceding dry conditions, and also these crops will require more rain to adequately sustain growth. Favourable conditions prevailed in northern Europe, the Atlantic region and in western and eastern Mediterranean countries.

Contents:

1. Agrometeorological overview
2. Remote sensing – observed canopy conditions
3. Pastures in Europe – regional monitoring
4. Country analysis and sowing updates
5. Crop yield forecasts
6. Atlas

Covers the period from 1 April to 10 May 2020




Crop	Yield (t/ha)				
	Avg 5yrs	April Bulletin	MARS 2020 forecasts	% Diff 20/5yrs	% Diff April
TOTAL CEREALS	5.60	5.44	5.39	-3.7	-0.9
Total Wheat	5.54	5.65	5.50	-0.6	-2.7
<i>soft wheat</i>	5.77	5.87	5.72	-0.9	-2.6
<i>durum wheat</i>	3.49	3.43	3.38	-2.9	-1.5
Total Barley	4.78	4.86	4.72	-1.1	-2.9
<i>spring barley</i>	4.02	4.03	4.05	+0.8	+0.5
<i>winter barley</i>	5.75	5.92	5.63	-2.0	-4.9
Grain maize	7.58	8.04	7.94	+4.8	-1.2
Rye	3.75	3.92	3.90	+4.0	-0.5
Triticale	4.04	4.18	4.15	+2.7	-0.7
Rape and turnip rape	3.08	3.14	2.95	-4.4	-6.1
Potato	32.5	34.2	33.9	+4.6	-0.8
Sugar beet	74.7	75.9	75.4	+0.9	-0.7
Sunflower	2.25	2.39	2.34	+4.4	-2.1

Issued: 15 May 2020


1. Agrometeorological overview

1.1. Areas of concern

AREAS OF CONCERN - EXTREME WEATHER EVENTS



AREAS OF CONCERN - CROPS



Large parts of western and central Europe experienced persistent precipitation deficits. In eastern France, eastern United Kingdom, the Benelux countries, northern half of Germany, and western Poland, as well as in southern Czechia, western Slovakia, Austria, eastern Italy, Slovenia, Hungary and Croatia, weather conditions were similar to those reported in the April Bulletin: low precipitation, high radiation levels and predominantly above-average temperatures. Some rainfall did take place (up to 60 mm in eastern France; generally less than 30 mm in the other aforementioned regions), but mostly in the last ten days of the review period. In general the precipitation amount was insufficient to replenish or sustain favourable soil moisture levels for the growth of winter and early planted spring crops (see dedicated sowing paragraphs in the April issue of the Bulletin). As a consequence, crops biomass accumulation slowed down with actual crop conditions generally worse than in 2019. Moreover, in most of these regions, little or no rain is forecast in the coming days, as winter cereals are approaching the sensitive flowering stage. In the case of rapeseed, the conditions of water stress already experienced partially coincided with the flowering and/or early grain filling stage. In eastern Romania and western Ukraine, the condition of crops worsened due to the ongoing drought and crops are

generally stunted and present signs of wilting and early leaf senescence.

For the sowing and emergence of late planted spring crops and summer crops (see sowing paragraphs of this Bulletin) in these regions, the concerns raised in the April Bulletin were mitigated. The limited amount of rainfall since the end of April was sufficient to facilitate the progress of sowing, and to allow for emergence (see section on sowing conditions); however, emerging crops often present uneven or patchy stands due to the preceding dry conditions, and also these crops will require more rainfall to adequately sustain growth.

A favourable rainfall surplus is observed in western and eastern Mediterranean countries. In the Maghreb region, rainfall occurred too late in the season to be beneficial for winter crop grain filling while unusually high temperatures in Tunisia impacted the grain filling of winter crops.

Note: Just after the review period, a marked cold-air intrusion caused night frosts mainly in Germany, the United Kingdom and the Benelux countries. Depending on the local climate conditions frost damages are possible for summer crops, flowering rapeseed, permanent crops, and vineyards.

1.2. Meteorological review (1 April – 10 May 2020)

Warmer than usual conditions were observed in most of France, and in north-eastern Spain, with daily mean temperature anomalies (with respect to the long-term average (LTA)) of between 2°C and 4°C.

Slightly warmer-than-usual conditions were observed across western and central Europe, the western Mediterranean region, and southern Sweden. Daily mean temperatures reached up to 2°C above the LTA. For over 7 days, the daily maximum temperatures were above 25°C in large areas of southern and eastern Spain, Po Valley (Italy), and the Balkans.



Slightly colder conditions than usual occurred in large regions of eastern Europe, the Mediterranean, Finland, Norway, and northern Sweden. The observed daily mean temperature anomalies did not exceed -2°C below the LTA. These areas received 11 to 15 cold days, with daily minimum temperatures of below 0 °C.



Wetter-than-usual conditions were reported in Spain and France, as well as in large regions of Greece, Turkey, central-northern Norway, Finland, and European Russia. In most of France and Spain, cumulated precipitation was 50% - 80% higher than the LTA during the review period. Anomalies of 40% above the LTA were recorded in the

other aforementioned areas.

Drier-than-usual conditions were recorded across a wide belt of central Europe, from Germany and Poland, to Czechia, Slovakia, and Austria, and further south in the Po Valley region of northern Italy, as well as in Slovenia, Croatia, and Hungary. The United Kingdom, Ireland, and southern Norway were also drier than usual. Furthermore, dry conditions prevailed locally in eastern European areas, such as Lithuania, western Belarus, eastern Romania, and southern Ukraine. In all of these regions, cumulated precipitation was mainly ranging between -80% and -50% below the LTA during the review period.

Focusing on a longer temporal perspective, since February, a three-month precipitation deficit has affected Hungary, Croatia, Slovenia, and the Po Valley region of Italy, as well as southern Russia and Ukraine. Over the past two months, the deficit has extended to a larger region across Europe, including Ireland, the United Kingdom, the Benelux countries, Germany, western and southern Poland, Slovakia, Austria, and eastern Ukraine. In contrast, the water supply from rainfall has been favourable in recent months for the Iberian Peninsula, south-western France, and Greece





1.3. Weather forecast (15 - 21 May 2020)

Weather conditions during the forecasted period will be mainly determined by an almost stationary pattern characterised by a cutoff cyclonic system in the western Mediterranean, and two troughs over the Atlantic and northern European Russia.

Warmer-than-usual conditions are predicted for south-eastern Europe and the eastern Mediterranean, with daily mean temperatures of over 4°C warmer than the long-term average (LTA), and up to 8°C above the LTA in Greece and Turkey. In these areas, several days with daily maximum temperatures of above 30°C are expected. Warmer conditions than usual are also forecast for southern Italy, as well as across some areas of western Europe: Portugal, the United Kingdom, and Ireland are expected to see daily mean temperature anomalies of between 2°C and 4°C.

Colder-than-usual conditions are forecast across central, eastern, and northern Europe, as well as in European Russia. Daily mean temperatures will stay around 2°C to 4°C below the LTA. In a large region across eastern Europe and Russia, as well as in Finland, central-northern Norway, and northern Sweden, the anomalies will be more pronounced, and are expected to be around 4° and 6°C below the LTA.



Accordingly, during the forecast period, an average temperature accumulation is expected in western Europe and south of the Alps, while it is forecast to be slightly



above average for the United Kingdom and Ireland, and it will remain below average in central-eastern and northern Europe. The temperature accumulation in south-eastern Europe is expected to be significantly above average.

Dry conditions with less than 5 mm of accumulated precipitation are expected in a wide belt spreading from Portugal and western Spain over most of France, the Benelux countries, Germany, and Poland, into southern Belarus, northern Ukraine, and western Russia. Also, Hungary, Slovakia, Turkey, and most of Greece, are expected to remain dry.

Wet conditions with accumulated precipitation of between **40 mm and 80 mm** (locally, 100 mm) are forecast in central Italy and south-eastern Europe, while around the Pyrenees and the Alpine regions, the Baltics, and Finland, up to 40 mm of precipitation are expected.


The **long-range weather forecast** for May, June, and July strongly suggest that warmer conditions than usual will be experienced in most of Europe, as well as very likely in the central Mediterranean, and in south-eastern Europe. These warmer conditions will likely go together with drier conditions than usual for south-eastern Europe.





2. Remote sensing – observed canopy conditions







Slowdown in biomass accumulation in central and eastern Europe





The map displays the differences between the fraction of Absorbed Photosynthetically Active Radiation (fAPAR) cumulated from 1 April to 10 May 2020, and the medium-term average (MTA, 2010-2019) for the same period. Positive anomalies (in green) reflect above average canopy density or early crop development, while negative anomalies (in red) reflect below average biomass accumulation, or late crop development.

In **Spain**, winter crops exhibit generally favourable biomass accumulation, and very advanced stages (e.g. *Castilla Y Leon*), thanks to persistent rainfall and warm weather. The flowering stage has almost ended, and the fAPAR curve displays a similar profile to 2016, a year which resulted in very good winter crop yields. In **southern Italy**, colder temperatures than usual until mid-April caused a slowdown in biomass accumulation (e.g. *Puglia*), and rainfall sustained an average crop development. Winter crops are flowering, and crops' conditions are mixed, still reflecting the unfavourable early spring conditions. In north-eastern **France**, the unfavourable start for spring crops, and the slowdown of winter crops' development, is reflected by the overall crops' biomass accumulation averaging lower than usual (light red on the map). In northern and north-western regions, winter crops' development remains slightly advanced, and vegetative growth is sustained, as a result of the favourable rainy and warm weather conditions which have taken place since the end of April. In **Germany**, the development of winter crops changed from very advance to slightly advanced, due to a combination of dropping temperatures at the beginning of April, and suboptimal soil moisture throughout the whole month. Crops' biomass accumulation slowed down significantly (e.g. *Thüringen*). Since the beginning of May, some rainfall has meant that crops have avoided early senescence. In **Poland**, crop conditions are similar to those observed in Germany, but the slowdown of biomass accumulation is more marked, due to very low soil moisture, which hampered crops' growth (e.g. *Lodzkie*). The winter crops' continue to be very advanced (green on the map), and the recent precipitation partially alleviated the dry conditions, but more rainfall is needed to further sustain crops' growth. In **the Baltics**, and **Finland**, the fAPAR displays very advanced stages; the biomass accumulation of winter crops is well sustained by the favourably distributed precipitation since 1 April. In central Europe (**Slovakia, the Czech Republic,**

Austria, and Hungary), the persistent dry conditions in April are similar to 2019 with respect to biomass accumulation, and crop development is advanced (e.g. *Niederösterreich*). In western regions of **Romania** and **Bulgaria**, the winter crop conditions are favourable, despite the unfavourable dry conditions throughout April. In eastern regions, the long-lasting deficit of precipitation led to an early senescence of winter crops, especially along the border with Moldova and Ukraine (e.g. *Sud-Est*); the rainfall of May only partially mitigated the expected yield loss. In **Ukraine**, the deficit of precipitation came to a stop in May. The previous lack of precipitation and the warm temperatures depleted topsoil moisture, but the very good soil moisture capacity maintained sufficient moisture for winter crops' growth (e.g. *Zaporis'ka*, see cloud effect in May), with the exception of the *Odeska* region, where crops are wilting. In **the United Kingdom**, the negative fAPAR anomaly is a consequence of a reduction of the area of winter cereals. This reduction was compensated by an increased area of spring cereals which are progressing well, and have benefitted from the precipitation in late April and early May (e.g. *Essex*). In **Turkey**, the delay of winter crops is not yet fully recovered, but the yield outlook is positive. In **Morocco**, the winter crop season ended with low yields).






3. Pastures in Europe – regional monitoring

Dry conditions limit recent grassland productivity

Pasture Productivity Index

Period of analysis: 11 April - 10 May 2020

Index based on MetOP-AVHRR fAPAR 10-day product.
Medium-term average (MTA) 2010-2019




The pasture productivity index (PPI) for the period of 11 April to 10 May is shown in the main map above. The following countries or regions predominantly experienced above average pasture productivity: Spain, Portugal¹, central Italy, Greece, Denmark, Sweden, Estonia, and Finland, where favourable rainfall and temperatures provided good biomass growth.

Though the map suggests predominantly above average pasture conditions across Europe, mainly triggered by the mild winter and spring temperatures, the appearance of yellow and reddish areas (when compared with the map published in the April issue of the Bulletin) indicates hampered productivity. The lack of an increase in pasture productivity over the spring months – albeit due to the mild conditions at a relatively high initial productivity level – is mainly to be attributed to the unusually dry conditions which have prevailed since mid-March in France (all areas except for the far west), Germany, Austria, Czechia, Poland, eastern Romania, Slovenia, Croatia, Slovakia, north-eastern Bulgaria, Latvia, Lithuania, and Ireland. Reports in the media suggest that the first cut is already compromised in some of these countries (eastern Romania, and certain areas of Poland). These two areas have been the most affected by the dry spell in the past months, and will require more regular rainfall to recover.

¹ The below-average PPI values in southern Portugal are attributed to excessive cloud cover, which hampered adequate satellite observations. Rainfall and temperature conditions in these regions have been favourable for pastures.

Germany - North


Reference period: 01 Apr to 10 May 2020



	BULLETIN ISSUE									
	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT		
RAINFALL	Light Green	Light Orange	Orange	White						
TEMPERATURE	Light Green	Light Green	Dark Green	White						
RADIATION	Light Green	Light Green	Dark Green	White						

France - East


Reference period: 01 Apr to 10 May 2020



	BULLETIN ISSUE									
	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT		
RAINFALL	Light Green	Light Green	Orange	White						
TEMPERATURE	Light Green	Light Green	Orange	White						
RADIATION	Light Orange	Light Green	Dark Green	White						

Finland and Baltic republics


Reference period: 01 Apr to 10 May 2020



	BULLETIN ISSUE									
	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT		
RAINFALL	Light Green	Light Green	Dark Green	White						
TEMPERATURE	Light Green	Light Green	Dark Green	White						
RADIATION	Light Orange	Light Green	Dark Green	White						

Romania - East


Reference period: 01 Apr to 10 May 2020



	BULLETIN ISSUE									
	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT		
RAINFALL	Light Orange	Light Orange	Orange	White						
TEMPERATURE	Light Green	Light Green	Dark Green	White						
RADIATION	Light Green	Light Green	Dark Green	White						

Poland


Reference period: 01 Apr to 10 May 2020



	BULLETIN ISSUE									
	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT		
RAINFALL	Light Green	Light Orange	Orange	White						
TEMPERATURE	Light Green	Light Green	Dark Green	White						
RADIATION	Light Green	Light Green	Dark Green	White						

Denmark and Sverige

Reference period: 01 Apr to 10 May 2020



	BULLETIN ISSUE									
	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT		
RAINFALL	Light Green	Light Green	Dark Green	White						
TEMPERATURE	Light Green	Light Green	Dark Green	White						
RADIATION	Light Green	Light Green	Dark Green	White						

4. Country analysis

4.1. Sowing conditions

Spring barley

Sowing almost completed, in dry conditions

In France, Ireland, the Benelux countries, Germany, and the United Kingdom, sowing was mainly concluded in good conditions, when soil moisture conditions were still adequate for sowing, as reported in the April issue of the Bulletin. In general, these countries experienced warmer temperatures than usual, which resulted favourably for emergence. However, the dry conditions created uneven emergence and gappy stands in some areas, particularly for the later-sown crops.

In Denmark (the EU's third largest producer, after Spain and France) and Sweden, sowing started later than usual, at the beginning of April, and the rainfall at the end of April and in early May improved soil conditions for emergence.

In Poland, where sowing was accomplished in early April, the rainfall at the end of April and the beginning of May

improved soil water conditions, but emergence remained uneven.

In Hungary, Romania, Czechia, Slovakia, and Croatia, sowing was concluded in dry conditions. Despite some recent rainfall, soil water deficit continues to prevail, causing loss to yield potentials.

Sowing is progressing well in the Baltic Sea region. Thanks to the mild weather conditions, the sowing campaign began earlier than usual, at the beginning of April in Latvia, and in mid-April in southern Finland. Cold weather, combined with some rainfall in May, slowed down sowing activities. Despite this, sowing continued within the normal window. Further north, where temperatures are still cold, sowing is expected to start after mid-May, which is usual for these regions.

Sugar beet and potatoes

Dry conditions hamper germination and emergence in the main producing regions

In general, weather conditions have been favourable for the timely sowing of sugar beet across Europe, resulting in the completion of sowing works within the optimal time window. However, dry conditions in April in the main EU sugar beet producing regions of France, Germany, Poland, the United Kingdom and the Benelux countries hampered germination and the emergence of seedlings, most markedly in Germany and Poland. The situation improved at the end of April, when scattered showers resulted in improved water conditions for topsoil. However, in many fields, germination was poor or uneven, and more rainfall is still needed in order to secure adequate early crop development. Additionally, in Poland, as with some other regions, emerging plants which have already been weakened by dry conditions could have been damaged by cold snaps, as well as strong pest pressure.

Rainfall deficit led to similar difficulties in eastern Austria, Hungary, western Slovakia, eastern Croatia, eastern Slovenia, and Romania.

In the United Kingdom, the situation looks more positive, as the early-sown sugar beet crops emerged well, but also there the dry April conditions resulted in an uneven emergence of sugar beet in many plots.

In northern Spain, the weather conditions were generally favourable for the development of sugar beet, whilst in Italy, dry conditions led to a suboptimal start to the season.

Similar challenges were faced for potatoes, of which the main production regions coincide with those of sugar beet. In several regions of central and north-western Europe, early-sown stands, were locally damaged due to frost events at the end of the review period. At the early stages of development potato crops normally recover well from such damage. In many regions, potato planting is still ongoing. Compared to initial plans, the area sown is likely to be somewhat reduced, due to a combination of the above-mentioned difficulties and the drastically reduced demand for fries.

Maize

Maize sowing and emergence still facing difficulties in several regions

The sowing campaign of maize started with some concerns about a possible drought and low soil moisture levels not allowing plants to emerge. While rainfall observed since the beginning of May has been beneficial in the most affected regions, more rain is still needed to sustain adequate crop establishment.

Maize sowing has almost completed in most of the main producing regions. However, difficulties are still observed and conditions are highly contrastful.

The long dry period observed in April allowed farmers to prepare the seedbeds despite the increasing difficulties (hard clods, dust) because of the dry upper soil layers.

In Romania, sowing is almost finished, and while the soil was dry at the start of the sowing campaign, conditions

have improved but some regions are still very dry. Fields that were partly dry at sowing now often present patchy stands. Ukraine and Bulgaria benefited from the recent rainfall.

In Hungary, Veneto in Italy, Germany, Lithuania and eastern Poland, lack of rainfall might result in very poor emergence of maize sown into dry soils and some of the fields might eventually not be sown.

In France, a larger area than last year was sown to maize, and sowings were slightly advanced. The French southwestern regions received a large rainfall surplus lastly and this will ensure at least the early development of maize. Spain also benefited from a large rainfall surplus.

Sunflowers

Problematic sowing campaign due to dry topsoils in several regions

In Romania, Hungary, and Croatia (respectively the EU's 1st, 3rd, and 9th largest sunflower producing countries), the sowing campaign began around 5 April; with a delay of 1 to 2 weeks due to lower-than-usual top-soil temperatures. Sowing progressed well from the second dekad of April until late April, thanks to scarce rainfall and near-average or above-average temperatures. However, the conditions for sprouting and early crop development gradually worsened, as soils became progressively drier, which may lead to uneven stands. Beneficial rainfall in early May somewhat mitigated the situation. However, more rainfall is needed to sustain adequate development. Similar sowing problems occurred further northwest, in Slovakia, Czechia, Austria, and Germany.

In Bulgaria and Greece (the EU's 2nd and 7th largest producers, respectively), the sowing campaign started, slightly delayed, during the second week of April. Since then, rainfall and thermal conditions have been favourable for making good progress, and for the

sprouting and early growth of sunflower plants.

In France (the EU's 4th largest sunflower producer), the sowing of sunflower plants took full swing in the second half of April. Rainfall was substantially above the LTA in all main sunflower producing regions, but caused only short temporal delays. Rain fell with moderate intensity, favouring water infiltration into the soil without water logging problems, and therefore providing excellent conditions for emergence, and crop establishment.

In Spain and Portugal (the EU's 5th and 13th largest producers, respectively), sizeable and well-distributed rainfall resulted in favourable conditions for soil preparation and sunflower sowing, which has now been mostly concluded. Also, in Italy (the EU's 6th largest producer), soil and weather conditions have been favourable for sunflower sowing and germination. In southern Italy, plants are now at the emergence stage. In northern Italy, the sowing ended more recently, and seeds are close to the emergence stage.

Soybeans

Soybean sowing campaign ongoing, but central Europe needs more rainfall

The soybean sowing campaign is reaching an end in France, Ukraine, Romania, and Bulgaria. In these countries, the recent rainfall will ensure the emergence of soybean. As is the case with maize, some of the concerns about a difficult start to the season because of the rainfall deficit are still valid. Most of central Europe would need at least some rainfall to ensure adequate emergence. This is

particularly the case in Hungary, north-western Austria, and eastern Poland. In some of the main soybean producing regions of Italy, the EU's largest soybean producing country, water availability for irrigation and linked canopy temperature control at the moment is not fully granted due to the low level of water reservoirs.

4.2. European Union

France

A challenging season and a particularly poor outlook for winter cereals





The succession of unfavourable weather conditions has provided a poor outlook for winter crops and spring cereals. Only summer crops are currently benefitting from a good start to the season.

Weather conditions during the review period were exceptionally warm in April, combined with a long period without rainfall in the north and east of France. Since the end of April, an exceptional rainfall surplus has been observed in southern France, but rainfall is still needed in northern and eastern France.

Throughout the review period, the dry and warm weather conditions hampered the early growth of spring barley as plants were exposed to water stress, and the same conditions favoured the development of aphids on winter wheat. Moreover, the record-warm conditions in April caused an acceleration of the phenological development of winter and spring cereals between the tillering and heading stages, thus diminishing the yield potential. This accelerated development is completing the large panel of unfavourable conditions observed since the start of the season (as exposed in earlier issues of the Bulletin): a wet autumn

hampering sowings and crop establishment, waterlogging hampering root growth, an exceptionally mild and wet winter favouring a high disease pressure, the delayed application of fertilisers at the start of a long dry period, as well as water stress. The recent rainfall limits further water stress, but increases disease pressure, particularly *septoria* for winter wheat, and *rhynchosporium* for winter barley. As the flowering stage for winter wheat is approaching, attention will be paid to wet conditions around flowering, which could favour the development of *fusarium*. Similar difficulties were experienced by winter rapeseed, and the yield forecasts for all winter crops are now well below the historical trend.

Conditions for summer crops have been mixed. Irrigated potato crops benefitted from a good start to the season, but rainfed potatoes and sugar beet have been exposed to water stress since their establishment. The main producing regions of grain maize, sunflower, and soybean, recently received a rainfall surplus, ensuring a good start to the season, which might be particularly valuable given the seasonal weather forecasts of a warm and dry summer.



Germany

Rainfall needed for an adequate crop development

April marked a record dry period throughout Germany, as most regions did not have a single day with significant rainfall. A huge radiation surplus is confirmed, and crops are now in urgent need of rainfall. Forecasts for winter crops are revised downwards.





Crop growth conditions during April were characterised by an almost complete absence of rain, and a very high level of sunshine. Some rainfall occurred during the last days of April and in the beginning of May, however this was not in the form of steady rain, but rather scattered showers. Consequently, the rain deficit remains huge, and concerns about a third consecutive dry year have not been alleviated. Soils in the north which were overly wet are now dried out, and a broad band with quickly depleting soil moisture is depicted from *Nordrhein-Westfalen*, and parts of *Schleswig-Holstein* towards the east (*Thüringen*, *Sachsen-Anhalt*, *Brandenburg*, *Sachsen*).

The phenological development of winter crops continues

to be advanced, but with a cooler weather forecast for Germany, a beneficial slowdown of growth is expected. Winter wheat has started heading, and precipitation is needed to sustain the yield potential. The forecast has been lowered, but it is still above last year's level.

Rapeseed in the west – under stressed water supply conditions, and having experienced a few cold spells, – is nearing the end of the flowering period. In *Bayern*, flowering started towards the end of April, and the flowers might be susceptible to the forecasted cold spells. The rapeseed forecast is lowered compared with the April Bulletin, and corresponds to the 5-year average.

Summer crops were sown, and emerged under dry conditions (albeit with difficulties on very light soils). However, their yield potential should not be reduced if water supply returns to average within the next few weeks.



Poland





Dry conditions reduce yield potential for winter and spring crops

During the review period, temperatures oscillated around the LTA. Brief cold spells occurred during the first two dekads of April, with temperature minima below -4°C . Exceptionally low rainfall in April (75% below LTA) resulted in very dry soils. However, the moderate amount of rainfall which has fallen since the end of April temporarily increased topsoil water content, partially relieving crops.

Rainfall at the end of April and the beginning of May brought critically needed water for winter crops which are entering the heading and flowering stages. The rain also improved topsoil water conditions for spring crops, and favoured the germination and early development of summer crops. Very dry conditions from mid-March to late April may negatively impact the yield potential for winter

and spring crops. This will largely depend on whether enough rain will fall in the coming weeks to sustain the adequate development of these crops. Cumulative radiation was above the LTA. Colder nights during the first dekads of April, with temperatures $\sim 0^{\circ}\text{C}$, slowed down crop development. Furthermore, frost events in April provided additional stress to the dry conditions for winter crops and emerging summer crops. In addition, sugar beet received strong pest pressures

Spring and summer crops' yield forecasts remain unchanged compared with last month, and follow the historical trend. Yield forecasts for winter wheat and barley have been slightly reduced, due to unfavourably dry conditions.



Ireland

Crops are progressing well despite dry conditions – but more rain is needed



Warmer temperatures advanced crop development, and recent rainfall improved soil moisture conditions, particularly for the early growth of spring crops. Spring and winter crops are forecast below last year's yield, but close to the five-year average.

Temperatures were above average, with cumulated radiation values higher than usual. Rainfall generally remained below average, but was more frequent and abundant in southern Ireland. Beneficial precipitation fell across the country from the end of April onward.

The sowing of spring cereals was mostly completed by mid-April; the warmer temperatures enabled crops to emerge quickly, but the lack of soil moisture created an uneven emergence. Spring crops benefitted from rainfall

at the end of April, and most crops are growing well. Sugar beet sowing is almost completed, and emergence is variable.

Winter wheat crops advanced in their development, and crop conditions are variable depending on the region and soil type, with better growing conditions in the southern areas, where rainfall has been more favourable. In some areas, the soil water content was depleted, and more rain is needed to maintain the yield potential. Generally, winter crops are around booting stage, the expected growth stage for this time of year. Winter oilseed rapeseed is at full flowering. Dry conditions are not considered to impact the yield at this stage, and our yield forecasts remain close to the 5-year average, but below the 2019 yield.



Spain and Portugal

Good crop growing conditions prevailing





Soil moisture levels recovered well in most regions, but still remain below optimum in the south. Nevertheless, the overall crop yield outlook is positive for both winter and summer crops.

Rainfall has been above average from 1 April onward in most of Spain and Portugal, and was around average in *Castilla La Mancha, Murcia, and Andalucia*. Most of the soil moisture on the Iberian Peninsula is now fully recovered from the long-lasting dry winter period (see soil moisture and cumulated water balance figures below). The south of Portugal and Spain received less rainfall in recent weeks than central regions; however, the situation in the south is still close to average conditions, and is remarkably better than it was two months ago. Temperatures across the Iberian Peninsula were warmer than average, and this was more pronounced in the north, thus continuing the accelerating effect of the warm winter on crop development (see Valladolid figure below).

Winter crop development is more advanced than usual across all of Spain and Portugal due to the cumulated effect of mild winter conditions, and the return of rainfall last month in the south, which has been beneficial for sustaining winter crop yields.

Regarding summer crops, the sowing of maize and sunflower was completed in good conditions, and initial soil water conditions have been favourable for their development. Limited water supply for irrigation may still be an issue in the coming months in southern Spain (*Andalucia, Guadiana, Guadalquivir*), where the accumulated water reserves amount to half, or less than half, of the 10-year average reservoir filling for May (source: www.embalses.net).

The yield outlook for both winter and summer crops is positive, and is in keeping with the forecast of the previous Bulletin.







Italy

Mixed conditions for winter crops

Winter wheat yield forecasts (soft and durum) are just below the 5-year average, but slightly above 2019, reflecting the mixed crops conditions: in central and southern regions, average to good growing conditions counterbalanced the adverse dry weather in the north. Maize and soybean were planted during April under dry conditions, while sunflower sowing recently concluded with favourable weather. Summer crops yield forecasts still reflect the trend value.

In north-western regions, May brought some rain, which partially mitigated the dry soil conditions. In northern and north-eastern regions, only 20 mm to 40 mm of rainfall has taken place since the beginning of April (20% to 50% compared with the LTA). During the review period, the warm weather (+2°C to +4°C compared with the LTA) accelerated the soil moisture depletion in the Po Valley, and in the north-eastern plain. Both areas present strong deficits in climatic water balance: -80 mm since 1 April, compared with a positive water balance of +20 mm in the LTA. The driest provinces (i.e. eastern Emilia Romagna, and southern Veneto) are relevant for soft wheat (50% of the

national production), maize (30%), and soybean (60%) production. Negative effects on yields due to the unfavourable conditions for soft wheat in those regions are likely to be compensated for at a national level by Piemonte and Lombardia's around average crop conditions. For maize and soybean, water availability for irrigation, and linked canopy temperature control is not currently fully granted, due to water reservoirs being low. In central regions, rainfall was almost seasonal, with 40 mm to 60 mm poured in just a few days at the beginning of May. Winter crops' development is delayed in Toscana, but is slightly advanced in Marche and Umbria, with fair biomass accumulation. Sunflower sowing proceeded favourably. In late April, all southern regions (with the exception of eastern Sicilia) received beneficial rainfall. In these regions, durum wheat and winter barley, now around the flowering stage, present mixed conditions, resulting from the occurrence of several suboptimal weather conditions along the campaign. However, the very negative expectations for early spring did not materialise.



Hungary

Recent rainfall was insufficient to relieve concerns

Dry weather conditions predominated until late April, when precipitation tendencies increased. However, this was insufficient for fully alleviating winter crops' soil moisture deficit. The persistently suboptimal soil moisture levels are increasingly concerning, as winter cereals are close to reaching the sensitive heading/flowering stages.





Temperatures fluctuated significantly, but were near-seasonal on average when considering the review period as a whole. A significant cold spell occurred in early April, and additional frost events took place in mid-April. The latter are likely to have caused only limited damage to the flowering rape-seed stands.

The very dry weather conditions from early March persisted throughout most of April. Rainfall became more frequent at the end of April, and in early May, but was insufficient to fully alleviate the rainfall deficit. Precipitation totals for the past 60 days were on average between 10 mm and 35 mm, which corresponds to 35% to 60% of the LTA. This period has been either the driest or second driest since 1979. The medium range weather

forecast predicts 5 mm to 15 mm of precipitation, which would be insufficient for significantly improving soil moisture levels, which are particularly low in eastern Hungary, and in some north-western regions.

The spring sowing campaign accelerated after mid-April, when soil temperatures reached adequate levels, but the quality of seed beds (dry, with many clods) was suboptimal. Recent rainfall is facilitating the sprouting, emergence, and early development of summer crops.

The development of winter crops remains advanced, due to the unusually high temperatures during winter and early spring. The flowering of rapeseed begun in mid-April, but water supply – crucial for yield formation – has been suboptimal during this period. The current situation is delicate for winter cereals, which are close to reaching the heading/flowering stage. Remote sensing images and crop model results still indicate near average or above average conditions, however, water stress is imminent, considering the unfavourable water supply and negative precipitation outlook. The yield forecasts for winter crops and summer crops were revised downwards.







Romania

Drought forming

A long lasting rain deficit resulted in serious crop water supply problems in eastern Romania (in Nord-Est, and more seriously in Sud-Est), and to a lesser extent, in the western regions along the Hungarian border.

Near average thermal conditions prevailed in eastern Romania, whereas the western half of the country was slightly colder than usual (1°C - 2°C), considering the review period as a whole. Daily mean temperatures fluctuated considerably around the LTA, and two significant frost events occurred around 1 and 15 April. Dry weather which had begun in March persisted in April. Precipitation totals in April reached only between 5 mm and 25 mm. The southern and eastern regions received hardly any precipitation. In early May, beneficial precipitation (10 mm to 40 mm) arrived throughout the country, partially alleviating the rain deficit. Soil moisture contents under winter crops are very low, primarily in the south and the east, where winter cereals

are reaching the start of the heading/flowering stage, which is cardinal for yield formation and production. So far, inadequate water supply most adversely impacted flowering rapeseed crops, but also affected the canopy expansion and biomass accumulation of winter cereals. The dry conditions allowed good progress of the spring sowing campaign, and the most recent rainfall improved soil moisture levels. However, the same conditions were unfavourable for the emergence and early growth of spring and summer crops, particularly in the cases of spring barley and sugar beet. The current medium range weather forecast suggests considerable rainfall in north-western Romania (*Vest, Nord-Vest, Centru, and Nord-Est*), but very little in southern regions. The yield forecasts for winter crops (particularly for rapeseed and winter wheat), as well as for spring crops, was revised considerably downward, and moderately downward for summer crops.



Bulgaria

Rainfall deficit in the north-eastern regions





Crop water supply was mostly adequate in Bulgaria, except in the north-eastern regions. Recent rainfall improved the situation even in the north-east, however, soil moisture levels remain low. Remote sensing indicators suggest a near average condition of winter crops.

Thermal conditions were near the average throughout the review period (1 April – 10 May), albeit with considerable fluctuations. The first dekad of April was predominantly colder than usual.

In northern and eastern Bulgaria, April was much drier than usual, with only 10 mm to 40 mm of rainfall (20% to 50% of the LTA). The most serious rainfall deficit occurred in north-eastern Bulgaria (*Severozitochen*, and *Severen Tsentralen*). However, the precipitation in early May mitigated the water supply problems in these regions. In contrast, the *Yuzhen Tsentralen* region received abundant precipitation (70 mm to 115 mm) during the review period, exceeding the LTA by 30% to 120%.

Topsoil conditions allowed adequate progress of the spring sowing campaign, except in southern-central Bulgaria, where it was temporarily hampered by wet weather.

Soil moisture levels under winter crops are generally near or above average, but very dry soil conditions prevail in north-eastern regions. The development of winter crops is advanced by 5-10 days. Remote sensing indicators show a near average biomass accumulation. In the north-eastern and eastern regions (where much of the rapeseed is cultivated), the break point in the course of the NDVI/fAPAR graphs (e.g. *Severen Tsentralen*) suggest that the situation has worsened from a good to an average level, due to inadequate water supply. In other regions, soil water supply is likely to be sufficient to meet the water demand of winter cereals in the near future. Our previous optimistic yield forecast for winter rapeseed was revised downward, whereas our forecast for other winter crops was revised upward, and is now at the trend level. .



Austria, Czechia, and Slovakia



Deepening soil moisture deficit

Less than 30 mm of rainfall was recorded since the beginning of April, which was not sufficient to replenish the soil moisture deficit in western Slovakia, eastern Austria, and eastern Czechia. The dry conditions are limiting winter crop growth, reducing the yield outlook to below the 5-year average. Recent rainfall slightly improved the emergence conditions for summer crops.

With temperatures up to 2°C above the LTA, the period from the beginning of April onwards has been warmer than usual in Austria and Czechia. Maximum temperatures reached up to 26 °C. After a completely dry first half of April, several rainfall episodes followed in mid-April and at the beginning of May with, however, very limited quantities. Since the beginning of April, the total rainfall recorded in Slovakia, eastern Austria, and eastern Czechia, was less than 30 mm, representing less than half of the

expected seasonal values in these regions.

The recent rainfall was insufficient to compensate for the pronounced soil moisture deficit in eastern Austria, western Slovakia, and eastern Czechia, limiting the growth of winter crops in these regions. Consequently, we revised the yield forecast of winter rapeseed downward. Due to the warm winter and spring so far, winter soft wheat is in the advanced development stage, approaching the sensitive flowering period; final yields will strongly depend on the rainfall in the coming weeks. Therefore, as with the April Bulletin, our forecast remains below the 5-year average. Also, the emergence of summer crops is hampered by the lack of moisture in many places, even though the recent rainfall has slightly improved surface soil moisture levels. It is still too early to understand the impact on summer crop yields, for which our current forecasts remain in line with the long-term trend.



Denmark and Sweden

Positive yield outlook maintained for winter crops

While a long dry period with a large radiation surplus was observed in April, weather conditions have returned to seasonal averages since the beginning of May. The yield forecasts for winter crops are maintained close to record levels.



While April was practically dry, substantial rainfall has taken place since the beginning of May, resulting in cumulative rainfall amounts for the entire review period of over 50% of the seasonal values in all regions. Temperatures were slightly above average in April, after which the temperature regime gradually returned to seasonal values. An exceptional radiation surplus was

observed throughout the review period.

Winter crops benefitted from sufficient soil moisture and the radiation surplus, while warm temperatures in April favoured crop growth. The yield forecast for winter cereals and winter rapeseed are maintained well above the historical trend, reflecting that many recent years have been characterised by an early start to spring.

Spring crops, and more particularly spring barley, which were largely sown after mid-March, benefitted from the rainfall at the beginning of May. The yield forecast for spring barley is maintained close to the trend, considering that the long dry period was not beneficial for crop

establishment. However, the recent rainfall can still improve crop conditions.



Finland, Lithuania, Latvia and Estonia



Spring sowings are progressing well

Winter crops are faring well in the Baltics and Finland. Although rainfall and cold weather slowed down spring sowing in Finland, sowing is still progressing within the expected window. Spring sowings are almost completed in the Baltic countries, where recent rainfall sustained the early growth stages.

Temperatures were warmer than usual in all agricultural regions until mid-April. After this point, temperatures became colder than usual, with minimum temperatures decreasing to below 0°C several times during the reporting period. Cumulative rainfall was below average in Lithuania and Latvia, and above average in the northern countries.

Winter crops are in good conditions, and soil moisture is adequate to sustain growth in all countries. Spring sowings progressed well in the Baltic countries. Spring crops benefitted from the rainfall at the beginning of May, which is confirmed by the adequate levels of soil water supply simulated by our model. Spring sowings started earlier than usual in Finland due to the warm weather and the small amount of snow in winter, but the cold weather and rainfall of May brought the work back to the usual schedule.

The winter crops' yield forecasts remain close to the five-year averages. For other crops, the yield forecast is still based on historical trends.



Belgium, Luxembourg and the Netherlands

More rain is needed to sustain a fair outlook



The rain since the end of April brought relief to winter crops, and improved soil conditions for the sowing and emergence of sugar beet, potatoes, and maize. Winter crops are in fair condition. However, soil moisture levels are still exceptionally low for May, and an above average rainfall will be needed to sustain good yields.

The exceptionally dry period from mid-March onwards ended on 28 April, and was followed by a rainy period until 4 May, after which it was largely dry again. Precipitation totals during the review period varied between 20 and 30 mm (35-55% of the LTA), with the lowest values in eastern Netherlands and north-eastern Belgium. Daily mean temperatures were predominantly above the LTA, but with large fluctuations. Minimum temperatures dropped to below 0°C for two to three days in the first half of April, and to just above 0°C for two to three days around 6 May. Radiation levels were exceptionally high.

In general, winter crops successfully withstood the unusually dry conditions until the end of April, and have benefitted from the improved conditions since then. Phenological development and biomass accumulation are above average. However, soil moisture levels are still exceptionally low, and, while little to no rain is forecast for the coming 10 days, above average rainfall of just the right amount will be needed to sustain good growth towards flowering and yield formation.

Conditions for recently sown crops improved with the recent rainfall. Sugar beet fields are uneven due to the dry conditions, and some fields were re-sown. The planting of potatoes and maize is well underway.

The yield forecasts for winter crops were revised slightly downwards, while the yield forecasts for summer crops are still based on historical trends.



Greece and Cyprus

Positive outlook for winter cereals and favourable conditions for summer crops



In Greece, wet spring conditions were beneficial to crops. Cereal biomass accumulation is generally above average. Timely fungal treatments helped to manage most of the biotic crop stress which resulted from the overly wet conditions in April. Summer crops are developing well, and have entered the post-emergence stage.

The review period was characterised by an above average rainfall which cumulated in eastern and central Greece, and by average precipitation levels along the Adriatic coast and in the Peloponnese. Most of the rain fell during the first two dekads of April, in events characterised by moderate intensity. Above average precipitation was also

observed in Cyprus, with fairly distributed events throughout the period. Temperatures have ranged from seasonal to slightly below average for both countries. No particular thermal anomalies were observed. Despite the overly wet conditions in spring, the situation for crops in Greece is generally favourable. The wet weather in March and April favoured fungal infections (i.e. *Septoria* spp.), which in most cases were treated on time, and thus have not damaged the winter crop yields. Winter wheat is proceeding well, and is now at grain filling stages. Summer crops, mainly cultivated in the northern regions of Greece, are growing under favourable conditions, and are in the first vegetative stages. The analyses of satellite indicators

confirm our positive expectations for both winter and summer crops. Biomass accumulation levels for rainfed and irrigated lands are above the medium term average

(MTA), and well above the previous season. Our forecasts are above the five-year average for winter crops, and follow the trend for summer crops.





Slovenia and Croatia

Dry conditions lowering winter crop yield outlook

The lack of rainfall since the beginning of the year has caused drought stress in many agricultural regions of Croatia and Slovenia; most markedly in eastern regions of both countries. The yield outlook for winter crops has been revised downwards. The lack of soil moisture is also hampering the emergence of summer crops.

Even though April began with cold weather, the period as a whole was slightly warmer than usual in Slovenia and western Croatia. Both countries received less than half of their usual rainfall during the review period, and this was most pronounced in central-western and eastern Croatia; eastern Croatia saw less than 40 mm of rainfall altogether. Adding to the precipitation deficit since the beginning of the year, soil moisture levels for winter crops are now well below the LTA in eastern Croatia, as well as in eastern Slovenia. The deficit is especially marked on sandy soils, and soils with little organic matter.

The freezing temperatures at the beginning of April caused no significant damage to winter crops. Currently, the main threat to winter crop growth, and the emergence of summer crops, comes from the prevailing drought conditions. Even though the cold weather anomaly at the end of March and the beginning of April slowed down the phenological development of winter crops, these are mainly in an advanced stage, due to the preceding warm weather. Winter soft wheat is gradually entering the flowering stage, which is highly sensitive to drought stress. Consequently, we revised downward the yield outlook for winter wheat, especially in Croatia. Similarly, the winter rapeseed outlook was also revised downward. For summer crop emergence, the lack of moisture in many places causes a problem. Nevertheless, it is still too early to quantify the impact on crop yields, and so our forecast remains around the long-term trend.



4.3. United Kingdom

United Kingdom

More rainfall is needed to maintain yield potentials



Above average temperatures accelerated the development of winter crops. April rainfall sustained winter and spring cereals, although more rain is needed to approach yield potential. Dry conditions created an uneven germination in spring barley and sugar beet.

Average temperatures were warmer than usual, with cumulated radiation above the LTA. The rainfall deficit continued throughout the review period, although moderate precipitation occurred since mid-April. In particular, this was Scotland's third driest period (1 April to 10 May) on record since 1979. This impacted spring barley germination, which is still not complete, and is uneven in many areas, particularly for crops which were sown later. Some farmers have irrigated spring barley fields.

Spring crops generally benefitted from the rainfall since mid-April, but more rain is needed to ensure optimal

growth. Sugar beet is emerging well with the early drillings, but at the end of April, sugar beet establishment was uneven in many fields. Potato planting is in full swing, and many growers have started irrigating, especially in the east.

Winter wheat growth stages vary according to the drilling date, but the September-drilled crops in the south of the UK have reached the flag leaf stage, while winter barley is at booting stage. Winter cereals are faring well, but soil moisture content decreased to below normal levels, according to our simulations. More rainfall is needed to maintain good yield potentials, particularly now that the crops are approaching critical growth stages. Due to the depleted soil water profiles, yield forecasts are revised slightly downwards for winter crops, but maintained to be close to the 5-year average for spring crops.



4.4. Black Sea Area

Ukraine

Wheat and barley yield forecasts lowered following the dry conditions





Despite the return of some rainfall since the beginning of May, the long dry period beforehand left a negative mark on the yield potential of winter cereals. Low temperatures and cold nights delayed the development of winter and spring crops.

The weather was particularly dry in April, with only a few oblasts (*Kyyivs'ka*, and *Vinnys'ka*) receiving substantial rainfall, whereas the rest of the country experienced no more than 50% of the usual rainfall. In the beginning of May, conditions changed, and since then, substantial rainfall has been observed in all oblasts except *Odes'ka*, which remained exposed to an exceptional drought. Temperatures were close to the seasonal average when considering the review period as a whole. However, daily amplitudes were large, and several night frost events were

reported.

Winter wheat was impacted by the dry conditions, and the yield forecast was revised downwards as the main producing regions (except *Vinnys'ka*) faced a water deficit. Winter barley in the main producing regions (*Odes'ka*, and *Mykolayivs'ka*) was also impacted by the water deficit, and the yield forecast is now below the trend. Spring barley development was hampered by the low soil moisture, delayed by the cold temperatures, and locally impacted by night frost.

While the sowing of summer crops was also hampered by the dry conditions and the cold (soil) temperatures, the precipitation observed since the beginning of May, combined with warmer temperatures, will finally allow the sowing campaign to be completed, and will ensure adequate emergence.




Turkey

Favourable winter crop season continues

Wheat and barley yield forecasting figures are above last year's, and the 5-year average. This is a result of very good crop conditions in south-eastern regions. Summer crop sowings are ongoing, and yield forecasts are based on trend analysis.

In *Konya* and *Kirikkale*, seasonal temperatures since 1 April favoured the normal development of winter crops. After a slow start to the season, crops recovered to normal stages, and an almost favourable biomass accumulation. In *Ankara* and *Kaysiri*, the temperature was slightly below average, and this slowed down crop growth, which had only recently reached average levels after an initial delay. In all aforementioned regions, favourable precipitation occurred at the beginning of May, replenishing soil moisture, and fully sustaining crop growth. In the southern Anatolian regions (*Konya*, and *Kirikkale*), winter crops are

almost at the heading stage, while in *Kaysiri*, winter crops are still delayed, and in full vegetative phase. In the Aegean regions (e.g *Adana*), where most of the maize is cultivated, a surplus of precipitation occurred during the spring, and water reservoirs for irrigation have been replenished. The sowing period has just begun, and maize yields will mostly depend on summer temperatures. In south-eastern regions (i.e. *Gaziantep*, *Sanliurfa*, and *Mardin*), favourable precipitation is the main driver for the very favourable biomass accumulation during winter crops' vegetative growth. Winter crops' flowering stage is now ending, and soil moisture is still sufficient to sustain grain filling. Yield expectations are very favourable. Water reservoirs are replenished, creating favourable expectations for the summer season as well.



4.5. European Russia and Belarus

European Russia

A tempered outlook for winter cereals; good progress of spring sowing campaign

Severe frost events in the first half of April are expected to have caused leaf and stem damages to winter cereals. Increasing soil moisture deficit has started to negatively affect the – so far very good – yield potentials in the main winter wheat producing regions in south-western Russia. The spring sowing campaign has begun early and is progressing well.





Most areas of European Russia largely experienced slightly colder thermal conditions than usual (0.5°C - 2°C < LTA) during the review period (1 April to 10 May), except for certain eastern and northern areas, where near-seasonal and somewhat warmer temperatures than usual prevailed. In the first half of April, extraordinary frost events, with daily minimum temperatures of -5°C to -10°C (locally down to -13°C) caused leaf and stem injuries of winter cereals, even in the southern regions.

Rainfall during the review period was abundant (40 mm – 90 mm) in the Central okrug (e.g. in the Black Earth region), and in the northern part of the Volga okrug. In contrast, in the southern and northern Caucasian okrugs, dry

conditions continued throughout, but the situation improved somewhat in early May, typically resulting in 10 mm - 40 mm for the review period as a whole (certain regions close to the Caucasus mountains received 50 mm - 100 mm of rainfall).

In southern Russia, after an excellent early start to spring, the growth of winter crops began to slow down due to the rapidly declining soil moisture levels. Nevertheless, crop model results and remote sensing images indicate still near average biomass accumulation, thanks to water reserves in deeper layers of the deep Chernozem soils. However, in the absence of very substantial rainfall, a drought situation can easily develop, as has been the case in the past two years.

The sowing of spring crops began early, and advanced significantly quicker than usual. Rain hampered the spring sowing campaign only in the Central and Volga okrugs. In southern regions, even the sowing of thermophile crops such as sunflower and maize has begun, and is progressing well.





Belarus

Average outlook despite a very dry April

During the review period, temperatures oscillated around the long term average (LTA). Frost conditions with temperature minima of lower than -3.5°C occurred briefly during the first two dekads of April. Cumulative precipitation in April was significantly below the LTA, at 65% below normal levels. This resulted in decreasing soil water levels in most of the country, especially in the south. Fortunately, precipitation events since the end of April alleviated dry topsoil conditions in most of the country, and brought some relief to crops.

Winter crops are generally in fair condition, developing at a normal pace, and currently in the tillering and stem elongation stages. Yield potentials of winter cereals may have been affected by the low soil moisture levels in April, especially in southern regions. Nevertheless, recent rainfall brought some relief to crops. The sowing campaign of spring and summer crops is close to being finalised under adequate soil water conditions for germination and emergence.

We maintain our yield forecasts based on historical trends.



4.6. Maghreb





Morocco, Algeria and Tunisia

Rain in western Maghreb was insufficient for crops to recover

The exceptional shortfall in precipitation since the beginning of the 2019-2020 growing season has negatively affected growth and development in most of the rain-fed cultivation in the Maghreb region. Late rainfall events in April and May were insufficient for crops to recover. Morocco is the worst-hit by the drought. Algeria is affected in the western regions, and Tunisia in the inland barley-producing regions.

In the Maghreb region, the review period (1 April to 10 May) has been characterised by rainfall events which mostly fell in western Maghreb. More precisely, rain fell across the northern belt to the east of *Tanger* (Morocco), and the west of *Nabeul* (Tunisia). Accumulated rainfall values ranged from 20 to 60 mm, which were 50-60% more than the LTA. Rain was 10 to 20 mm below average in central and southern Morocco, and at average levels in central Tunisia. Precipitation arrived too late to stimulate crop recovery in Morocco and Eastern Algeria, but was beneficial for soft and durum wheat in northern Tunisia.

Crops in Morocco have almost completed the phenological cycle, which was accelerated during February and March in response to drought conditions. By contrast, biomass levels are above average in eastern Algeria (e.g. *Guelma*, *Tebessa*) and in the northern littoral regions of Tunisia (e.g. *Bizerte*, and *Jendouba*), where grain-filling in April occurred under favourable soil moisture conditions. For the regions of *Le Kef*, *Siliana*, *Kasserine*, and *Kairouan* in central Tunisia, remote sensing indicators suggest below average biomass accumulation during grain-filling. This trend was already observed during flowering in March, and is a consequence of the scarce precipitation in these regions during spring, but also of the exposure to above-average temperatures, including a heatwave during the review period, in early May. The outlook for barley production in these regions deteriorated. Our forecasts generally reaffirm the negative expectations for cereals in Morocco, remain close to average for Algeria, and are revised downward for barley in Tunisia.





5. Crop yield forecast

Country	TOTAL WHEAT (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	5.54	5.77	5.50	-0.6	-4.7
AT	5.45	5.77	5.17	-5.1	-10
BE	8.47	9.23	9.01	+6.4	-2.4
BG	4.93	5.14	5.10	+3.5	-0.8
CY	-	-	-	-	-
CZ	5.93	5.73	5.76	-3.0	+0.4
DE	7.50	7.40	7.59	+1.2	+2.6
DK	7.66	8.19	7.88	+2.9	-3.8
EE	3.97	5.07	3.98	+0.3	-21
EL	2.62	2.65	2.79	+6.3	+5.2
ES	3.14	3.04	3.15	+0.4	+3.5
FI	3.91	4.56	3.89	-0.4	-15
FR	6.99	7.84	6.86	-1.9	-13
HR	5.54	5.53	4.97	-10	-10
HU	5.27	5.28	4.96	-5.9	-6.0
IE	9.84	9.99	9.84	-0.1	-1.6
IT	3.94	3.75	3.87	-1.9	+3.1
LT	4.48	4.29	4.59	+2.4	+7.1
LU	5.78	6.01	5.95	+3.0	-1.0
LV	4.49	4.81	4.53	+0.8	-5.9
MT	-	-	-	-	-
NL	8.90	9.44	8.90	+0.0	-5.7
PL	4.49	4.39	4.71	+4.9	+7.3
PT	2.19	2.23	2.46	+12	+10
RO	4.44	4.80	4.31	-3.0	-10
SE	6.55	7.40	7.34	+12	-0.8
SI	4.99	5.23	5.03	+0.7	-3.9
SK	5.15	4.81	4.83	-6.2	+0.5

UK	8.34	8.94	8.01	-4.0	-10
----	------	------	------	------	-----

Country	TOTAL BARLEY (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	4.78	4.99	4.72	-1.1	-5.4
AT	5.67	6.07	5.70	+0.6	-6.0
BE	8.06	8.56	8.68	+7.6	+1.4
BG	4.40	4.90	4.89	+11	-0.1
CY	1.67	2.70	1.55	-7.4	-43
CZ	5.33	5.38	5.26	-1.4	-2.1
DE	6.66	6.78	6.52	-2.1	-3.9
DK	5.60	6.29	5.84	+4.2	-7.2
EE	3.45	4.09	3.71	+7.4	-9.4
EL	2.60	2.71	2.76	+5.8	+1.8
ES	2.93	2.76	2.93	-0.1	+6.4
FI	3.72	4.23	3.72	+0.0	-12
FR	6.45	7.08	6.11	-5.3	-14
HR	4.79	5.18	4.92	+2.9	-5.0
HU	5.07	5.54	5.14	+1.5	-7.1
IE	8.01	8.66	7.86	-1.8	-9.2
IT	4.00	4.05	3.94	-1.4	-2.7
LT	3.36	3.37	3.35	-0.4	-0.5
LU	-	-	-	-	-
LV	3.19	3.43	3.33	+4.7	-2.7
MT	-	-	-	-	-
NL	6.44	6.51	6.75	+4.8	+3.7
PL	3.56	3.46	3.60	+1.2	+4.0
PT	2.39	2.64	2.38	-0.4	-10
RO	4.04	4.44	3.98	-1.7	-11
SE	4.68	5.31	4.96	+6.0	-6.5
SI	4.65	4.85	4.49	-3.4	-7.4
SK	4.66	4.81	4.57	-2.0	-4.9

UK	6.27	6.92	6.14	-2.2	-11
----	------	------	------	------	-----





Country	SOFT WHEAT (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	5.77	6.00	5.72	-0.9	-4.7
AT	5.52	5.83	5.22	-5.4	-11
BE	8.47	9.23	9.01	+6.4	-2.4
BG	4.93	5.14	5.10	+3.5	-0.8
CY	-	-	-	-	-
CZ	5.93	5.73	5.76	-3.0	+0.4
DE	7.52	7.42	7.62	+1.3	+2.6
DK	7.66	8.19	7.88	+2.9	-3.8
EE	3.97	5.07	3.98	+0.3	-21
EL	2.70	2.77	2.88	+6.5	+4.0
ES	3.23	3.09	3.25	+0.6	+5.0
FI	3.91	4.56	3.89	-0.4	-15
FR	7.11	7.92	6.95	-2.2	-12
HR	5.54	5.53	4.97	-10	-10
HU	5.29	5.32	4.98	-5.9	-6.3
IE	9.84	9.99	9.84	-0.1	-1.6
IT	5.41	5.14	5.28	-2.6	+2.6
LT	4.48	4.29	4.59	+2.4	+7.1
LU	5.78	6.01	5.95	+3.0	-1.0
LV	4.49	4.81	4.53	+0.8	-5.9
MT	-	-	-	-	-
NL	8.90	9.44	8.90	+0.0	-5.7
PL	4.49	4.39	4.71	+4.9	+7.3
PT	2.19	2.23	2.46	+12	+10
RO	4.44	4.80	4.31	-3.0	-10
SE	6.55	7.40	7.34	+12	-0.8
SI	4.99	5.23	5.03	+0.7	-3.9
SK	5.22	4.87	4.88	-6.4	+0.3

UK	8.34	8.94	8.01	-4.0	-10
----	------	------	------	------	-----

Country	DURUM WHEAT (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	3.49	3.47	3.38	-2.9	-2.4
AT	4.58	4.83	4.46	-2.7	-7.7
BE	-	-	-	-	-
BG	-	-	-	-	-
CY	-	-	-	-	-
CZ	-	-	-	-	-
DE	5.06	4.92	4.99	-1.5	+1.4
DK	-	-	-	-	-
EE	-	-	-	-	-
EL	2.59	2.61	2.76	+6.4	+5.7
ES	2.73	2.76	2.60	-4.6	-5.7
FI	-	-	-	-	-
FR	5.29	6.28	5.13	-2.9	-18
HR	-	-	-	-	-
HU	4.71	4.34	4.49	-4.6	+3.5
IE	-	-	-	-	-
IT	3.34	3.15	3.26	-2.6	+3.3
LT	-	-	-	-	-
LU	-	-	-	-	-
LV	-	-	-	-	-
MT	-	-	-	-	-
NL	-	-	-	-	-
PL	-	-	-	-	-
PT	-	-	-	-	-
RO	-	-	-	-	-
SE	-	-	-	-	-
SI	-	-	-	-	-
SK	4.53	4.29	4.31	-4.9	+0.5

UK	-	-	-	-	-
----	---	---	---	---	---





Country	SPRING BARLEY (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	4.02	4.13	4.05	+0.8	-1.8
AT	4.44	4.17	4.37	-1.5	+4.8
BE	-	-	-	-	-
BG	-	-	-	-	-
CY	-	-	-	-	-
CZ	5.18	5.07	5.12	-1.1	+1.0
DE	5.21	5.12	5.37	+3.1	+4.9
DK	5.44	6.14	5.71	+5.0	-7.0
EE	3.45	4.09	3.71	+7.4	-9.4
EL	-	-	-	-	-
ES	3.01	2.80	2.96	-1.8	+5.4
FI	3.72	4.23	3.72	+0.0	-12
FR	6.22	7.04	6.03	-3.1	-14
HR	-	-	-	-	-
HU	3.96	4.56	3.63	-8.4	-21
IE	7.27	8.00	7.35	+1.2	-8.1
IT	-	-	-	-	-
LT	3.34	3.29	3.26	-2.2	-0.9
LU	-	-	-	-	-
LV	3.19	3.43	3.33	+4.7	-2.7
MT	-	-	-	-	-
NL	6.44	6.51	6.75	+4.8	+3.7
PL	3.37	3.21	3.40	+0.9	+6.0
PT	-	-	-	-	-
RO	2.73	2.81	2.40	-12	-14
SE	4.61	5.19	4.84	+5.1	-6.7
SI	-	-	-	-	-
SK	4.45	4.57	4.29	-3.5	-6.2

UK	5.74	6.34	5.73	-0.2	-9.6
----	------	------	------	------	------

Country	WINTER BARLEY (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	5.75	6.11	5.63	-2.0	-7.8
AT	6.34	6.74	6.34	+0.0	-5.8
BE	8.06	8.56	8.68	+7.6	+1.4
BG	4.40	4.90	4.89	+11	-0.1
CY	1.67	2.70	1.55	-7.4	-43
CZ	5.68	5.98	5.54	-2.5	-7.4
DE	7.09	7.22	6.83	-3.6	-5.4
DK	6.47	7.03	6.63	+2.4	-5.7
EE	-	-	-	-	-
EL	2.60	2.71	2.76	+5.8	+1.8
ES	2.41	2.31	2.73	+13	+18
FI	-	-	-	-	-
FR	6.54	7.09	6.15	-5.9	-13
HR	4.79	5.18	4.92	+2.9	-5.0
HU	5.28	5.65	5.39	+2.2	-4.5
IE	9.25	9.42	9.22	-0.4	-2.1
IT	4.00	4.05	3.94	-1.4	-2.7
LT	3.90	3.89	4.04	+3.4	+3.8
LU	-	-	-	-	-
LV	-	-	-	-	-
MT	-	-	-	-	-
NL	-	-	-	-	-
PL	4.26	4.31	4.29	+0.9	-0.3
PT	2.39	2.64	2.38	-0.4	-10
RO	4.48	4.91	4.49	+0.2	-8.6
SE	5.91	6.85	6.61	+12	-3.4
SI	4.65	4.85	4.49	-3.4	-7.4
SK	5.21	5.29	5.06	-2.8	-4.5

UK	7.16	7.84	6.87	-4.1	-12
----	------	------	------	------	-----





Country	GRAIN MAIZE (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	7.58	7.90	7.94	+4.8	+0.6
AT	10.1	10.4	10.3	+2.0	-1.3
BE	10.4	10.5	11.0	+5.8	+5.7
BG	6.44	6.96	7.47	+16	+7.3
CY	-	-	-	-	-
CZ	7.30	8.29	7.63	+4.5	-8.0
DE	9.21	8.81	9.61	+4.3	+9.1
DK	-	-	-	-	-
EE	-	-	-	-	-
EL	10.3	10.6	11.0	+7.0	+3.9
ES	11.6	11.8	11.5	-0.4	-2.3
FI	-	-	-	-	-
FR	8.85	8.58	8.85	+0.0	+3.1
HR	7.87	9.01	8.22	+4.5	-8.8
HU	7.50	8.05	7.38	-1.7	-8.4
IE	-	-	-	-	-
IT	10.2	10.0	10.5	+3.0	+4.5
LT	6.39	7.67	6.84	+6.9	-11
LU	-	-	-	-	-
LV	-	-	-	-	-
MT	-	-	-	-	-
NL	9.78	9.75	10.7	+9.7	+10
PL	6.09	5.61	6.37	+4.7	+14
PT	8.52	8.98	8.82	+3.5	-1.9
RO	5.52	6.52	6.25	+13	-4.1
SE	-	-	-	-	-
SI	8.85	9.27	8.89	+0.4	-4.1
SK	6.94	7.39	7.52	+8.4	+1.8

UK	-	-	-	-	-
----	---	---	---	---	---

Country	RYE (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	3.75	3.87	3.90	+4.0	+0.8
AT	4.43	4.60	4.59	+3.6	-0.3
BE	-	-	-	-	-
BG	-	-	-	-	-
CY	-	-	-	-	-
CZ	4.93	5.06	5.00	+1.4	-1.3
DE	5.12	5.24	5.25	+2.5	+0.2
DK	5.89	6.10	6.11	+3.9	+0.3
EE	3.61	4.12	3.60	-0.3	-13
EL	1.72	1.86	1.90	+10	+2.1
ES	2.10	1.82	2.04	-2.8	+13
FI	3.79	4.82	3.72	-1.7	-23
FR	4.52	4.77	4.55	+0.7	-4.5
HR	-	-	-	-	-
HU	3.14	3.37	3.12	-0.5	-7.3
IE	-	-	-	-	-
IT	-	-	-	-	-
LT	2.51	2.63	2.55	+1.7	-2.9
LU	-	-	-	-	-
LV	4.14	4.43	4.21	+1.6	-5.0
MT	-	-	-	-	-
NL	-	-	-	-	-
PL	2.77	2.72	2.89	+4.2	+6.0
PT	0.95	1.06	1.04	+9.2	-2.2
RO	-	-	-	-	-
SE	6.17	6.76	6.58	+6.5	-2.8
SI	-	-	-	-	-
SK	3.44	3.44	3.56	+3.6	+3.5

UK	2.22	2.38	2.26	+1.6	-5.3
----	------	------	------	------	------





Country	TRITICALE (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	4.04	4.06	4.15	+2.7	+2.2
AT	5.36	5.49	5.48	+2.2	-0.2
BE	-	-	-	-	-
BG	2.96	2.84	3.02	+2.1	+6.5
CY	-	-	-	-	-
CZ	4.79	4.93	4.66	-2.9	-5.5
DE	6.01	6.13	6.16	+2.4	+0.5
DK	-	-	-	-	-
EE	-	-	-	-	-
EL	2.11	2.24	2.33	+10	+3.7
ES	2.35	2.32	2.70	+15	+16
FI	-	-	-	-	-
FR	5.04	5.44	5.03	-0.2	-7.6
HR	3.93	3.98	3.79	-3.6	-4.7
HU	3.96	3.95	3.92	-1.1	-0.9
IE	-	-	-	-	-
IT	-	-	-	-	-
LT	3.36	3.29	3.26	-2.9	-0.9
LU	-	-	-	-	-
LV	-	-	-	-	-
MT	-	-	-	-	-
NL	-	-	-	-	-
PL	3.55	3.49	3.68	+3.6	+5.5
PT	1.67	1.47	1.54	-7.9	+5.2
RO	3.90	4.12	3.88	-0.5	-5.7
SE	5.58	6.36	6.21	+11	-2.3
SI	-	-	-	-	-
SK	-	-	-	-	-

UK	4.33	4.48	4.10	-5.2	-8.5
----	------	------	------	------	------

Country	RAPE AND TURNIP RAPE (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	3.08	2.96	2.95	-4.4	-0.4
AT	3.08	2.98	2.92	-5.4	-2.2
BE	3.87	3.52	4.00	+3.5	+14
BG	2.73	2.67	2.78	+1.6	+4.1
CY	-	-	-	-	-
CZ	3.26	3.05	3.00	-8.0	-1.7
DE	3.39	3.30	3.39	+0.0	+2.7
DK	3.92	4.40	4.36	+11	-1.0
EE	2.14	2.64	2.20	+3.1	-17
EL	-	-	-	-	-
ES	2.10	2.13	2.08	-1.2	-2.5
FI	1.49	1.33	1.49	-0.6	+12
FR	3.33	3.11	3.03	-8.9	-2.5
HR	2.78	2.50	2.79	+0.2	+12
HU	3.08	2.97	2.86	-6.9	-3.6
IE	-	-	-	-	-
IT	2.60	2.66	2.64	+1.4	-0.6
LT	2.73	2.85	2.66	-2.5	-6.7
LU	-	-	-	-	-
LV	2.75	2.93	2.85	+3.7	-2.7
MT	-	-	-	-	-
NL	-	-	-	-	-
PL	2.77	2.73	2.77	-0.1	+1.3
PT	-	-	-	-	-
RO	2.57	2.04	2.08	-19	+2.2
SE	3.18	3.62	3.53	+11	-2.6
SI	-	-	-	-	-
SK	3.02	2.84	2.73	-9.4	-3.8

UK	3.55	3.31	3.46	-2.7	+4.5
----	------	------	------	------	------





Country	SUGAR BEETS (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	74.7	NA	75.4	+0.9	NA
AT	70.8	70.5	71.8	+1.5	+1.9
BE	84.9	88.2	85.5	+0.6	-3.1
BG	-	-	-	-	-
CY	-	-	-	-	-
CZ	62.7	61.8	61.7	-1.5	-0.2
DE	73.6	72.7	77.2	+4.9	+6.1
DK	69.9	80.7	67.5	-3.5	-16
EE	-	-	-	-	-
EL	-	-	-	-	-
ES	91.0	96.7	89.8	-1.3	-7.2
FI	37.9	47.6	40.9	+7.9	-14
FR	87.0	84.7	88.4	+1.7	+4.4
HR	60.1	52.8	56.5	-6.0	+7.0
HU	62.3	58.4	59.6	-4.4	+2.0
IE	-	-	-	-	-
IT	64.1	NA	62.2	-3.0	NA
LT	59.3	71.0	56.0	-5.7	-21
LU	-	-	-	-	-
LV	-	-	-	-	-
MT	-	-	-	-	-
NL	83.1	83.9	89.0	+7.1	+6.0
PL	61.8	NA	57.0	-7.8	NA
PT	-	-	-	-	-
RO	38.4	31.1	36.9	-3.7	+19
SE	63.6	74.0	63.6	+0.0	-14
SI	-	-	-	-	-
SK	59.7	57.6	58.7	-1.7	+1.8

UK	70.4	69.0	74.2	+5.3	+7.5
----	------	------	------	------	------

Country	POTATO (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	32.5	NA	33.9	+4.6	NA
AT	30.3	31.3	32.0	+5.6	+2.2
BE	41.0	41.1	45.2	+10	+10
BG	16.5	14.5	16.0	-2.9	+10
CY	-	-	-	-	-
CZ	26.9	27.2	27.7	+3.0	+1.9
DE	41.8	39.0	44.1	+5.5	+13
DK	41.0	42.5	42.1	+2.6	-0.9
EE	-	-	-	-	-
EL	27.6	27.7	27.6	-0.1	-0.5
ES	31.5	33.1	31.4	-0.3	-5.1
FI	27.4	28.9	28.6	+4.3	-1.1
FR	41.2	41.4	43.1	+4.5	+4.0
HR	-	-	-	-	-
HU	24.2	25.0	25.0	+3.2	-0.1
IE	-	-	-	-	-
IT	28.3	NA	27.7	-2.3	NA
LT	15.8	18.1	16.3	+3.3	-9.8
LU	-	-	-	-	-
LV	19.6	NA	19.8	+1.3	NA
MT	-	-	-	-	-
NL	42.0	42.0	44.0	+4.6	+4.6
PL	25.7	NA	25.1	-2.4	NA
PT	20.8	22.7	22.3	+7.0	-1.8
RO	15.6	14.8	16.3	+4.3	+10
SE	34.2	35.8	33.2	-2.9	-7.3
SI	-	-	-	-	-
SK	-	-	-	-	-

UK	40.5	36.5	41.8	+3.2	+15
----	------	------	------	------	-----





Country	SUNFLOWER (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	2.25	2.32	2.34	+4.4	+1.2
AT	2.68	3.00	2.70	+0.9	-10
BE	-	-	-	-	-
BG	2.28	2.31	2.55	+12	+10
CY	-	-	-	-	-
CZ	2.43	2.44	2.41	-0.7	-1.0
DE	2.02	2.04	2.04	+1.1	-0.2
DK	-	-	-	-	-
EE	-	-	-	-	-
EL	2.59	2.80	2.65	+2.1	-5.6
ES	1.15	1.12	1.20	+3.7	+6.3
FI	-	-	-	-	-
FR	2.28	2.19	2.36	+3.5	+7.5
HR	2.88	2.89	2.96	+2.9	+2.3
HU	2.88	3.00	2.97	+3.0	-1.1
IE	-	-	-	-	-
IT	2.37	2.47	2.40	+1.0	-3.0
LT	-	-	-	-	-
LU	-	-	-	-	-
LV	-	-	-	-	-
MT	-	-	-	-	-
NL	-	-	-	-	-
PL	-	-	-	-	-
PT	1.49	1.76	1.93	+30	+9.9
RO	2.47	2.64	2.51	+1.4	-5.1
SE	-	-	-	-	-
SI	-	-	-	-	-
SK	2.66	2.64	2.76	+3.7	+4.6

UK	-	-	-	-	-
----	---	---	---	---	---

Country	SOYBEAN (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
EU	2.90	2.96	2.95	+1.5	-0.4
AT	2.86	3.11	2.78	-2.8	-11
BE	-	-	-	-	-
BG	1.36	2.00	2.01	+48	+0.4
CY	-	-	-	-	-
CZ	2.10	2.27	2.28	+8.5	+0.4
DE	-	-	-	-	-
DK	-	-	-	-	-
EE	-	-	-	-	-
EL	-	-	-	-	-
ES	-	-	-	-	-
FI	-	-	-	-	-
FR	2.61	2.39	2.72	+4.3	+14
HR	2.74	2.90	2.89	+5.4	-0.5
HU	2.60	2.78	2.57	-1.0	-7.6
IE	-	-	-	-	-
IT	3.61	3.66	3.64	+0.6	-0.7
LT	-	-	-	-	-
LU	-	-	-	-	-
LV	-	-	-	-	-
MT	-	-	-	-	-
NL	-	-	-	-	-
PL	-	-	-	-	-
PT	-	-	-	-	-
RO	2.39	2.55	2.45	+2.5	-4.0
SE	-	-	-	-	-
SI	-	-	-	-	-
SK	2.22	2.46	2.56	+15	+4.0

UK	-	-	-	-	-
----	---	---	---	---	---



Country	WHEAT (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
BY	3.51	3.83	3.81	+8.5	-0.5
DZ	1.57	NA	1.55	-1.3	NA
MA	1.89	1.54	1.56	-18	+1.2
TN	1.77	NA	1.86	+4.8	NA
TR	2.78	2.78	2.94	+5.7	+5.9
UA	4.01	4.16	4.09	+1.7	-1.8

Country	GRAIN MAIZE (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
BY	5.86	6.00	5.76	-1.9	-4.1
DZ	-	-	-	-	-
MA	-	-	-	-	-
TN	-	-	-	-	-
TR	9.40	9.40	9.64	+2.5	+2.5
UA	6.59	7.19	7.39	+12	+2.7

Country	BARLEY (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
BY	3.08	3.50	3.38	+9.9	-3.4
DZ	1.28	NA	1.23	-4.1	NA
MA	1.27	NA	1.01	-21	NA
TN	0.83	NA	0.74	-10	NA
TR	2.70	2.64	2.77	+2.7	+5.1
UA	3.19	3.42	3.28	+2.8	-4.2

Country	SOYBEAN (t/ha)				
	Avg 5yrs	2019	MARS 2020 forecasts	%20/5yrs	%20/19
BY	-	-	-	-	-
DZ	-	-	-	-	-
MA	-	-	-	-	-
TN	-	-	-	-	-
TR	4.33	4.25	4.58	+5.7	+7.7
UA	2.19	2.29	2.37	+8.3	+3.4

Note: Yields are forecast for crops with more than 10000 ha per country with sufficiently long and coherent yield time series (for rice more than 1000 ha per country)

Sources: EU 2015-2020 data come from DG AGRICULTURE short term Outlook data (dated April 2020, received on 30/04/2020), EUROSTAT Eurobase (last update: 04/05/2020) and EES (last update: 15/11/2017)

Non-EU 2015-2019 data come from USDA, DSASI-MADR Algeria , INRA Maroc, CNCT Tunisie, Turkish Statistical Institute (TurkStat), EUROSTAT Eurobase (last update: 04/05/2020), State Statistics Service of Ukraine, FAO and PSD-online







2020 yields come from MARS CROP YIELD FORECASTING SYSTEM (output up to 10/05/2020)



EU aggregate after 01/02/2020 is reported.

NA = Data not available.





6. Atlas







Temperature regime







Precipitation











Climatic water balance





Crop development stages and precocity









Relative soil moisture



Maximum temperature around crops development



Precipitation around crops development



JRC MARS Bulletins 2020

Date	Publication	Reference
27 Jan	Agromet analysis	Vol. 28 No 1
17 Feb	Agromet analysis	Vol. 28 No 2
23 Mar	Agromet analysis, yield forecast	Vol. 28 No 3
27 Apr	Agromet analysis, remote sensing, pasture analysis, sowing conditions, yield forecast	Vol. 28 No 4
18 May	Agromet analysis, remote sensing, pasture analysis, sowing update, yield forecast	Vol. 28 No 5
15 Jun	Agromet analysis, remote sensing, pasture analysis, rice analysis, yield forecast	Vol. 28 No 6
27 Jul	Agromet analysis, remote sensing, pasture analysis, harvesting conditions, yield forecast	Vol. 28 No 7
24 Aug	Agromet analysis, remote sensing, pasture update, harvesting update, yield forecast	Vol. 28 No 8
14 Sep	Agromet analysis, remote sensing, pasture analysis, rice analysis, harvesting update, yield forecast,	Vol. 28 No 9
26 Oct	Agromet analysis, pasture update, sowing conditions, harvesting update, yield forecast	Vol. 28 No 10
23 Nov	Agromet analysis, sowing update, harvesting update	Vol. 28 No 11
14 Dec	Agromet analysis	Vol. 28 No 12

Mission statement: As the science and knowledge service of the European Commission, the Joint Research Centre's mission is to support EU policies with independent evidence throughout the whole policy cycle.

The current [JRC MARS Bulletin – Crop monitoring in Europe](#) is a JRC–European Commission publication from MARS4CAST (JRC Unit D5 – Directorate for Sustainable Resources)

[JRC MARS Bulletins](#) are available at <https://ec.europa.eu/jrc/en/mars/bulletins>

Analysis and reports

B. Baruth, S. Bassu, A. Bussay, A. Ceglar, I. Cerrani, Y. Chemin, P. De Palma, D. Fumagalli, R. Lecerf, G. Manfron, L. Nisini, L. Panarello, G. Ronchetti, L. Seguini, A. Toreti, M. van den Berg, M. van der Velde, Z. Zajac, A. Zucchini

Reporting support

SeproTec, I. Biavetti, G. Mulhern

Edition

M. van den Berg, B. Baruth, M. van der Velde, S. Niemeyer

Data production

MARS4CAST (JRC Unit D5), WENR (NL), MeteoGroup (NL), VITO (BE)

Contact

JRC D5/MARS4CAST
JRCMARSBULLETIN@ec.europa.eu

Legal notice

Neither the European Commission nor any person acting on behalf of the Commission is responsible for the use, which might be made of this publication.

Disclaimer

The geographic borders are purely a graphical representation and are only intended to be indicative. The boundaries do not necessarily reflect the official European Commission position.

Technical note

The long-term average (LTA) used within this Bulletin as a reference is based on an archive of data covering 1979-2019.

[Pre-press version](#). This is a pre-press version of the JRC MARS Bulletin, which, after final editing will be formally published by the Publications Office of the European Union.

In terms of content, both versions are identical.

